


1. *If $\phi(k) = A(a - |k|)$, $|k| \leq a$, and 0 elsewhere. Where a is a positive parameter and A is a normalization factor to be found.

(a) Find the Fourier transform for $\phi(k)$

(b) Calculate the uncertainties Δx and Δp and check whether they satisfy the uncertainty principle.

Soln



$$\int_{-\infty}^{\infty} \phi(k) \cdot \phi^*(k) \cdot dk = 1 \quad \left\{ \because \phi(k) \text{ is normalized} \right\}$$

$$\Rightarrow \int_{-a}^0 A^2 (a+k)^2 \cdot dk + \int_0^a A^2 (a-k)^2 \cdot dk = 1$$

$$\Rightarrow \frac{A^2}{3} (a^3) + \frac{A^2}{3} (a^3) = 1$$

$$\Rightarrow \frac{2A^2}{3} a^3 = 1$$

$$A = \pm \sqrt{\frac{3}{2} a^3}$$

(Usually taken as +ve one)

$$(a) \quad f(n) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \phi(k) e^{ikn} \cdot dk = \frac{1}{\sqrt{2\pi}} \left[\int_{-a}^0 (k+a) e^{ikn} \cdot dk + \int_0^a (a-k) e^{ikn} \cdot dk \right]$$

Subst. $k = -k$

$$\hookrightarrow \int_a^0 (-k+a) e^{-ikn} (-dk) = \int_0^a (a-k) e^{-ikn} dk$$


$$= \frac{1}{\sqrt{2\pi}} \left[\int_0^a (a-k) (e^{ikn} + e^{-ikn}) \right] \cdot dk$$

$$= \frac{1}{\sqrt{2\pi}} \int_0^a 2(a-k) \cos kn \cdot dk$$

$$= \sqrt{\frac{2}{\pi}} \left[\frac{(a-k) \sin kn}{n} \Big|_0^a + \int_0^a \frac{\sin kn}{n} \cdot dk \right]$$

$$= \sqrt{\frac{2}{\pi}} \left[\frac{(1 - \cos an)}{n^2} \right]$$





$$p = \hbar k \Rightarrow \Delta p = \hbar \Delta k$$

$$\Delta x \cdot \Delta k = \frac{4\pi}{a} \cdot 2a = 8\pi = \Delta x \cdot \frac{\Delta p}{\hbar}$$

$$\Delta x \cdot \Delta p = 8\pi \hbar$$

2. A wave packet is of the form $f(x) = \cos^2\left(\frac{x}{2}\right)$ (for $-\pi \leq x \leq \pi$) and $f(x) = 0$ elsewhere

(a) Plot $f(x)$ versus x .


(b) Calculate the Fourier transform of $f(x)$, i.e. $g(k) = \int_{-\infty}^{+\infty} f(x) e^{-ikx} dx$?

(c) At what value of k , $|g(k)|$ attains its maximum value?

(d) Calculate the value(s) of k where the function $g(k)$ has its first zero.

(e) Considering the first zero(s) of both the functions $f(x)$ and $g(k)$ to define their spreads

(i.e. Δx and Δk), calculate the uncertainty product $\Delta x \cdot \Delta k$.



$$\cos^2\left(\frac{u}{2}\right) = \frac{1}{2} (1 + \cos u)$$

$$\begin{aligned} \text{Sol}^n \quad (b) \quad g(k) &= \int_{-\pi}^{\pi} \cos^2\left(\frac{u}{2}\right) e^{-iku} \cdot du = \int_0^{\pi} \cos^2 \frac{u}{2} \left(e^{-iku} + e^{iku} \right) \cdot du \\ &= \int_0^{\pi} \frac{(1 + \cos u)}{2} (2 \cos ku) \cdot du \\ &= \int_0^{\pi} \cos ku + \frac{1}{2} (\cos(k+1)u + \cos(k-1)u) \cdot du \\ &= \frac{\sin k\pi}{k} + \frac{\sin(k+1)\pi}{2(k+1)} + \frac{\sin(k-1)\pi}{2(k-1)} \\ &= \frac{\sin k\pi}{k} - \frac{1}{2} \left[\frac{\sin k\pi}{(k+1)} + \frac{\sin k\pi}{(k-1)} \right] \\ &= \sin k\pi \left[\frac{(k^2-1) - (k^2)}{k(k^2-1)} \right] \\ &= \frac{-\sin k\pi}{k(k^2-1)} = \frac{\sin(k\pi)}{k-k^3} = g(k) \end{aligned}$$


$$(c) \quad |g(k)| = \left| \frac{\sin(k\pi)}{k-k^3} \right| = \frac{|\sin k\pi|}{|k-k^3|}$$

$$(c) |g(k)| = \left| \frac{\sin(k\pi)}{k-k^3} \right| = \frac{|\sin(k\pi)|}{|k-k^3|} \quad \text{looks like this}$$


$$\lim_{k \rightarrow 0} \frac{-\sin k\pi}{k(k-1)(k+1)} = +\pi$$

$$\lim_{k \rightarrow 1} \frac{-\sin k\pi}{k(k-1)(k+1)} = \lim_{k \rightarrow 1} \frac{\sin(k-1)\pi}{k(k-1)(k+1)} = +\frac{\pi}{2}$$

$$\lim_{k \rightarrow -1} \frac{-\sin k\pi}{k(k-1)(k+1)} = \lim_{k \rightarrow -1} \frac{+\sin(k+1)\pi}{k(k-1)(k+1)} = +\frac{\pi}{2}$$



Since $|k-k^3|$ is



(d) Zeros when $\sin k\pi = 0$ ($k \neq 0, 1, -1$)
 $\Rightarrow k = \pm 2, \pm 3, \dots$

$$\Delta k = 4$$

$$\Delta n, \Delta k = 8\pi$$

particle in a box

3. *A wave function $\psi(x)$ is defined such that $\psi(x) = \sqrt{2/L} \sin(\pi x/L)$ for $0 \leq x \leq L$ and $\psi(x) = 0$ otherwise.

(a) Writing $\psi(x) = \int_{-\infty}^{\infty} a(k) e^{ikx} dk$, find $a(k)$.

(b) What is the amplitude of the plane wave of wavelength L constituting $\psi(x)$?

$$\psi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} a(k) e^{ikx} dk$$

$$a(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \psi(x) e^{-ikx} dx$$

Solⁿ $a(k) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \psi(x) e^{-ikx} dx$

$$\begin{aligned} &= \frac{1}{2\pi} \int_0^L \sqrt{\frac{2}{L}} \sin\left(\frac{\pi x}{L}\right) e^{-ikx} dx = \frac{1}{2\pi} \int_0^L \sqrt{\frac{2}{L}} \sin\left(\frac{\pi x}{L}\right) (\cos kx - i \sin kx) dx \\ &= \frac{1}{2\pi} \sqrt{\frac{2}{L}} \int_0^L \frac{1}{2} (\sin\left(\frac{\pi}{L} + k\right)x) + \frac{1}{2} (\sin\left(\frac{\pi}{L} - k\right)x) dx \\ &\quad + \frac{1}{2\pi} \sqrt{\frac{2}{L}} (-i) \int_0^L \frac{1}{2} (\cos\left(\frac{\pi}{L} - k\right)x - \cos\left(\frac{\pi}{L} + k\right)x) dx \end{aligned}$$

$$= \frac{1}{2\pi} \sqrt{\frac{2}{L}} \frac{1}{2} \left[\frac{1 + \cos kL}{\frac{\pi}{L} + k} + \frac{1 + \cos kL}{\frac{\pi}{L} - k} \right] - \frac{i}{2\pi} \sqrt{\frac{2}{L}} \cdot \frac{1}{2} \left[\frac{\sin kL}{\frac{\pi}{L} - k} + \frac{\sin kL}{\frac{\pi}{L} + k} \right]$$

$$= \frac{1}{2\pi} \sqrt{\frac{2}{L}} \frac{1}{2} \left[\frac{1 + \cos kL}{\frac{\pi}{L} + k} + \frac{1 + \cos kL}{\frac{\pi}{L} - k} \right] - i \int \frac{(\sin kL)(2\pi)}{\dots}$$

$$= \frac{1}{2\pi} \sqrt{\frac{L}{2}} \left[\frac{(1 + \cos kL)(2\pi)}{(\pi^2 - k^2 L^2)} \right] - i \left[\frac{(\sin kL)(2\pi)}{(\pi^2 - k^2 L^2)} \right]$$

$$= \sqrt{\frac{L}{2}} \frac{1}{(\pi^2 - k^2 L^2)} \left[(1 + \cos kL) - i (\sin kL) \right]$$

(b) $f(n) \longleftrightarrow \sum_k a(k) e^{ikn}$

$\uparrow$ corresponding amplitude
 $\downarrow$ plane wave with wave number k

$$a\left(k = \frac{2\pi}{L}\right) = \sqrt{\frac{L}{2}} \left(\frac{1}{-3\pi^2} \right) (2)$$